

# Identification of relevant regions on structural and nonstructural proteins of *Zika virus* for vaccine and diagnostic test development: an *in silico* approach

E. A. Salvador<sup>1</sup>, G. A. Pires de Souza<sup>1</sup>, L. C. Cotta Malaquias<sup>1</sup>, T. Wang<sup>2</sup> and L. F. Leomil Coelho<sup>1</sup>

1) Institute of Biomedical Sciences, Department of Microbiology and Immunology, Federal University of Alfenas, Minas Gerais, Brazil and 2) Department of Microbiology & Immunology, Department of Pathology, Center for Biodefense and Emerging Infectious Diseases, University of Texas Medical Branch, Galveston, TX, USA

## Abstract

*Zika virus* (ZIKV) is an arbovirus belonging to the *Flaviviridae* family and the genus *Flavivirus*. Infection with ZIKV causes a mild, self-limiting febrile illness called Zika fever. However, ZIKV infection has been recently associated with microcephaly and Guillain-Barré syndrome. Vaccines for the disease are a high priority of World Health Organization. Several studies are currently being conducted to develop a vaccine against ZIKV, but until now there is no licensed ZIKV vaccine. This study used a novel immunoinformatics approach to identify potential T-cell immunogenic epitopes present in the structural and nonstructural proteins of ZIKV. Fourteen T-cell candidate epitopes were identified on ZIKV structural and nonstructural proteins: pr<sup>36–50</sup>; C<sup>61–75</sup>; C<sup>103–117</sup>; E<sup>374–382</sup>; E<sup>477–491</sup>; NS2a<sup>90–104</sup>; NS2a<sup>174–188</sup>; NS2a<sup>179–193</sup>; NS2a<sup>190–204</sup>; NS2a<sup>195–209</sup>; NS2a<sup>200–214</sup>; NS3<sup>175–189</sup>; and NS4a<sup>82–96</sup>; NS4a<sup>99–113</sup>. Among these epitopes, only E<sup>374–382</sup> is a human leukocyte antigen (HLA) type I restricted epitope. All identified epitopes showed a low similarity with other important flaviviruses but had a high conservation rate among the ZIKV strains and a high population coverage rate. Therefore, these predicted T-cell epitopes are potential candidates targets for development of vaccines to prevent ZIKV infection.

© 2019 Published by Elsevier Ltd.

**Keywords:** Diagnostic test, Immunoinformatic, T-cell epitope, vaccine, *Zika virus*

**Original Submission:** 28 October 2018; **Revised Submission:** 11 January 2019; **Accepted:** 15 January 2019

**Article published online:** 31 January 2019

**Corresponding author:** L. F. Leomil Coelho, Laboratório de Vacinas, Instituto de Ciências Biomédicas, Universidade Federal de Alfenas, Rua Gabriel Monteiro, 700 Centro, Alfenas Minas Gerais, 37130-001, Brazil.

**E-mail:** [luiz.coelho@unifal-mg.edu.br](mailto:luiz.coelho@unifal-mg.edu.br)

## Introduction

*Zika virus* (ZIKV) is an arbovirus transmitted in urban cycles to humans by the bite of infected female mosquitoes of the *Aedes* genus, mainly *Aedes aegypti*. ZIKV is a positive-sense, single-strand RNA virus classified in the genus *Flavivirus*, family *Flaviviridae* [1–3]. The virus genome contains one open reading frame, which is translated into a large polyprotein that should

be cleaved by viral and cellular proteases in order to generate three structural proteins (capsid, membrane and envelope proteins) and seven nonstructural proteins (NS1, NS2a, NS2b, NS3, NS4a, NS4b and NS5) [4].

ZIKV was first isolated in 1947 in Uganda [5], and six decades after its discovery, ZIKV caused an epidemic in the Federated States of Micronesia [6]. In 2013 the virus was introduced to Brazil, and after this event, a major pandemic in the Americas was observed [7–9]. The spread of ZIKV in the Americas was associated with the appearance of severe adverse outcomes, such as congenital Zika syndrome and Guillain-Barré syndrome [10–12]. Additionally the possibility of sexual transmission has intensified efforts to develop a vaccine for ZIKV [13,14]. Until now several approaches have been used to develop ZIKV vaccine, such as DNA and messenger RNA vaccines, inactivated ZIKV strains and recombinant proteins [15–17]. In general these vaccines are immunogenic and able to

induce protection in mice and nonhuman primates after challenge. However, vaccines developed using these approaches have yet to complete phase 2 clinical trials [18,19]. In addition, the success of preclinical test of these vaccines—the development of an effective ZIKV vaccine—could be complicated by previous immunity against other flaviviruses such as *Dengue virus* (DENV) [20–22]. Additionally, the antibody-dependent enhancement (ADE) phenomenon could confound the development of any flavivirus vaccine [23].

Traditionally vaccines have been developed by isolating one or more antigenic components from a given pathogen and checking whether these components are able of inducing a protective immune response. Although this approach has presented a series of successes throughout the history of vaccinology, the knowledge generated with the new genomic, transcriptomic and proteomic technologies has contributed to the advancement of vaccinology and the production of more effective and safe vaccines [24–26]. One of the most promising approaches is the use of computational tools that allow the identification of genes in the genome of different pathogens encoding proteins with antigenic potential. This approach, called reverse vaccinology (Rvac), is a direct consequence of new ‘omic’ technologies and is currently recognized as a promising technique for the theoretical determination of proteins or peptides that have a potential for induction of an immune response. The use of Rvac could reduce time and cost related to development of new vaccines [27–33]. Therefore, Rvac can find important peptide sequences that can be used in the development of new vaccines [33].

To overcome the confounding effects of cross-reactivity between DENV and ZIKV, the potential influence of a previous anti-flavivirus immunity on the outcome of ZIKV infection, the difficulty that this cross-reactivity implies in the development of vaccines and diagnostic tests of high sensitivity and specificity, we used the strategy of Rvac to find conserved, exclusive and potential immunogenic regions present in the structural and nonstructural proteins of ZIKV. These regions could be used as a guide to develop new vaccines and diagnostic technologies for the disease.

## Materials and methods

### T-cell epitope prediction

T-cell epitopes of all structural and nonstructural protein sequences of a Brazilian ZIKV strain (PE243/2015; GenBank accession no. ANC90426.1) were predicted by using the TepiTool tool available in the Immune Epitope Database (IEDB) Analysis Resource (<http://tools.iedb.org/tepitool>) [34]. MHC-I and MHC-II are highly polymorphic molecules with very

different allele specificity. Therefore, all predictions were performed to cover polymorphic loci using the most representative alleles in each class. The NetMCHpan method was used for MHC-I binding predictions and the NetMCHpanII for MHC-II binding predictions. A binding affinity threshold of 500 nM was selected as the cutoff for MHC-I binding and a 1000 nM cutoff for MHC-II binding, according to parameters recommended by TepiTool. The number of binding human leukocyte antigen (HLA) I and HLA-II alleles was recorded for each epitope.

### Epitope specificity analysis

The selected epitopes for ZIKV were compared for similarity with seven viruses belonging to family *Flaviviridae*, genus *Flavivirus* (Table 1). The full polyprotein sequences of these viruses were retrieved from GenBank (<http://www.ncbi.nlm.nih.gov/genbank>) and submitted to epitope conservation analysis using the Epitope Conservancy Analysis tool available online at the IEDB Analysis Resource (<http://tools.iedb.org/conservancy>) [35]. Data from previous studies have shown that epitopes with a high degree of similarity in amino acid sequences and three-dimensional structures could be recognized by cross-reaction antibodies [36]. Structural bioinformatics analysis showed cross-reactivity between allergenic proteins available with sequence identity >60% [37]. Therefore, only epitopes that showed a similarity of <60% with all other flaviviruses were considered to be specific for ZIKV and used for downstream analysis. Therefore, epitopes with similarity  $\geq 60\%$  with one or more virus were excluded from downstream analysis.

### Selection of immunogenic epitopes

The epitopes with low identity to other flaviviruses and a potential to interact with four or more HLA-I or HLA-II alleles were assessed for their antigenicity through the Vaxijen server, version 2.0 (<http://www.ddg-pharmfac.net/vaxijen/Vaxijen/Vaxijen.html>). For antigenicity analysis, a threshold of  $\geq 0.5$  was adopted, which corresponds to an accuracy of 87% for viruses [38]. The epitopes with values of  $\geq 0.5$  were submitted


**TABLE 1. Genomic sequences of flaviviruses included in this study**

Vector	Name	GenBank accession no.
Tick	Omsk hemorrhagic fever virus	NP_878909.1
	Tick-borne encephalitis virus	ANN44512.1
Mosquito	Dengue virus serotype 1	AGN94865.1
	Dengue virus serotype 2	AGE89225.1
	Dengue virus serotype 3	AHG23238.1
	Dengue virus serotype 4	ANK35834.1
	Japanese encephalitis virus	ANH21067.1
	St. Louis encephalitis virus	AIW82235.1
	West Nile virus	ALA10710.1
	Yellow fever virus	AHF35033.1
	Zika virus	ANC90426.1

to a conservation analysis of amino acid sequences among South American and African ZIKV isolates. Nucleotide and amino acid sequences of 684 ZIKV isolates were obtained from GenBank (<https://www.ncbi.nlm.nih.gov/genbank>) for comparative analyses using Epitope Conservancy Analysis tool available at the IEDB Analysis Resource (<http://tools.iedb.org/conservancy>). All epitopes presenting antigenic potential were also evaluated for theoretical determination of toxicity by using ToxinPred. This tool has an accuracy of 90% (<http://crdd.osdd.net/raghava/>

[toxinpred](#)) [39] and is widely used in the context of Rvac [40–42].

The selected epitopes were submitted to population coverage determination based on the group of alleles with which each epitope can interact according to the initial prediction using TepiTool. The calculation was performed using the Population Coverage Analysis tool of the IEDB [43]. Finally, the predicted epitopes on the M, E and NS1 proteins were evaluated for the solvent accessible surface (SAS) in the



**FIG. 1.** Schematic representation of entire *in silico* approach for predictions and selection of T-cell epitopes on structural and nonstructural Zika virus proteins.

corresponding protein using UCSF Chimera software version 1.12 and the 3-D cryostructure of E and M protein trimer (5IRE) and NS1 dimer protein (5GS6) deposited on the Protein Data Bank server.

## Results

As illustrated in Fig. 1, a nine-step immunoinformatic approach was used to identify T-cell epitopes with the potential to contribute to the development of vaccine candidates and diagnostic tests for ZIKV. Using TepiTool, 1457 epitopes were predicted within ZIKV polyprotein. Among these predicted epitopes, 944 (64.79%) were predicted for binding to HLA-I and 513 for HLA-II (31.21%). A total of 311 epitopes were identified from structural proteins and 1146 epitopes from

**TABLE 2. T-cell epitopes predicted from Zika virus structural and nonstructural proteins by TepiTool HLA-I and HLA-II prediction programme**


Protein	No. of predicted epitopes	
	HLA-I	HLA-II
C	40	19
PR	21	74
M	31	12
E	104	10
NS1	75	49
NS2a	96	35
NS2b	35	18
NS3	151	93
NS4a	43	20
2k	8	2
NS4b	96	41
NS5	244	140
Total	944	513

HLA, human leukocyte antigen.

nonstructural proteins (Table 2). Conservation analysis of predicted epitopes showed that 1019 (88.92%) had a higher similarity degree with other flaviviruses ( $\geq 60\%$  similarity) (Supplementary Table S1). Therefore, only 438 predicted epitopes (11.08%) had a similarity of  $< 60\%$  with structural and nonstructural proteins from other flaviviruses. Among these epitopes, only 187 (42.69%) showed a potential to interact with four or more HLA-I or HLA-II alleles (Fig. 2 and Supplementary Table S2). The data showed that HLA-II predicted epitopes are probably able to bind more alleles than the HLA-I predicted epitopes.


The immunogenicity potential of all 187 epitopes was evaluated using the Vaxijen server. The results showed that 87 epitopes (46.52%) were not predicted to be immunogenic. Thus, the number of candidates was reduced to 100 peptides with a range of antigenic potential score of between 0.5011 and 1.9567 (Fig. 3). The theoretical determination of toxicity was performed using ToxinPred, and only the epitope pr<sup>43-51</sup> showed a potential to trigger toxicity *in vivo* (Supplementary Table S3). Therefore, this epitope was excluded from the next analysis.

The 99 selected epitopes were submitted to a new conservation analysis using ten more sequences of DENV serotype 1, ten of DENV serotype 2, ten of DENV serotype 3, ten of DENV serotype 4, ten of Japanese encephalitis virus, four of Omsk hemorrhagic fever virus, nine of tick-borne encephalitis virus, four of West Nile virus, 16 of St Louis encephalitis virus and eight of yellow fever virus (Supplementary Table S4). Only eight epitopes presented a similarity of  $\geq 60\%$  with other flavivirus sequences; these were excluded from downstream analysis. These immunogenic epitopes were submitted to an analysis of conservation using 684 strains of ZIKV (Supplementary Table S5). The analysis showed a high degree of



**FIG. 2.** Number of HLA-I and HLA-II binding alleles by each predicted epitopes. Predicted epitopes with similarity less than 60% with structural and nonstructural proteins from other flaviviruses and theoretical number of binding HLA-I or HLA-II for each epitope are shown. Dotted line represents cutoff value (four or more binding HLA-I or HLA-II alleles). Only epitopes above cutoff were submitted to antigenicity calculation through Vaxijen server version 2.0 (<http://www.ddg-pharmfac.net/vaxijen/Vaxijen/Vaxijen.html>). HLA, human leukocyte antigen.

**FIG. 3.** Antigenicity prediction score computed by Vaxijen server. Antigenicity score was calculated by Vaxijen server version 2.0 (<http://www.ddg-pharmfac.net/vaxijen/Vaxijen/Vaxijen.html>). Dotted line represents cutoff value (> 0.5). Only epitopes above cutoff were submitted to conservation analysis of amino acid sequence between different South American and African Zika virus strains.



identity of all 100 epitopes among the ZIKV strains (90.89–100%) (Fig. 4). The evaluation of the population coverage based on the HLA alleles that each epitope can theoretically interact with varied from 17.68% and 59.52% for the epitopes predicted for HLA-I and from 59.39% and 99.97% for the epitopes for HLA-II. The HLA-II–predicted epitopes showed the highest coverage ( $\geq 50\%$ ) if compared to HLA-I–predicted epitopes ( $\leq 50\%$ ) (Fig. 5). The 91 selected epitopes and their properties are listed in Supplementary Table S6.


To gain insight into the spatial localization of the selected epitopes on E, M and NS1 proteins, these epitopes were located in the cryostructure of their respective proteins. The localization showed that all epitopes were exposed on the protein structures (Supplementary Fig. S1). The SAS calculation showed a high value of all epitopes ( $1471.71 \pm 309.2$ ) (Table 3).

In order to identify the most promising candidates for the development of future vaccines and/or to create diagnostic


tests to identify ZIKV, the selected epitopes were ranked. Further selection of epitopes with conservation rate  $\geq 95\%$  among the ZIKV strains, population coverage  $> 50\%$ , antigenic score  $\geq 1.0$  and SAS  $\geq 1050 \text{ \AA}^2$  (in this case, epitopes for M, E and NS1 proteins) resulted in a final list of 14 candidates epitopes (Table 4). These epitopes were located on C (2/14.3%), E (2/14.3%), prM (1/7.1%), NS2a (6/42.9%), NS3 (1/7.1%) and NS4a (2/14.3%) proteins. Among these epitopes, only epitope E<sup>374–382</sup> was a HLA-I–predicted epitope.

### Discussion

ZIKV is a mosquito-borne *Flavivirus* belonging to the *Flaviviridae* family [5]. It has been classified as an emerging pathogen because of its fast spread across the Americas as well as its association with cases of Guillain-Barré syndrome and prenatal



**FIG. 4.** Amino acid sequence identity of predicted epitopes among Zika virus (ZIKV) isolates. Peptide conservation analysis was performed by Epitope Conservancy Analysis tool at Immune Epitope Database (IEDB) (<http://tools.iedb.org/conservancy>). Six hundred eighty-four ZIKV isolates were included in this analysis. (A) Amino acid sequence identity of predicted epitopes for HLA-I alleles. (B) Amino acid sequence identity of predicted epitopes for HLA-II alleles. HLA, human leukocyte antigen.



**FIG. 5.** Classification of predicted epitopes based on population coverage rate. Selected epitopes were submitted to population coverage determination based on group of alleles that each epitope can interact according to initial prediction using TepiTool. Calculation was performed using Population Coverage Analysis tool of Immune Epitope Database (IEDB).

**TABLE 3.** Predicted epitopes for proteins E, M and NSI and calculation of SAS

Protein	Position	Epitope	HLA class	SAS (Å <sup>2</sup> )
M	2	VTLPSHSTR	I	1014.30
	60	KVIYLVML	I	968.19
	51	WLLGSSTSQKVIVLV	II	1430.80
	56	STSQKVIYLVMLLI	II	1386.61
E	46	TTVSNMAEV	I	1187.39
	164	RAKVEITPNSPRAEA	II	1886.30
	198	FSDLYYLTMINNKHVWL	II	1742.05
	269	LAGALEAEMDGAKGR	II	1569.30
	276	EMDGAKGRLSSGHLK	II	1776.53
	344	QMAVDMQTLTPVGRL	II	1376.53
	374	MMLELDPPF	I	1217.23
	477	LNTKNGSISLMCLAL	II	1252.47
NSI	58	SSVSRMENIMWRSVE	II	1794.59
	84	VQLTVVYVGSVKINPMW	II	2026.50
	125	RAAKTNSNF	I	1446.91

HLA, human leukocyte antigen; SAS, solvent-accessible surface.

microcephaly in ZIKV-endemic regions [46–48]. Therefore, there is an urgent need for vaccines against ZIKV infection [49–52]. Different immunoinformatics approaches aiming to identify the immunogenic peptides in the context of Rvac have been reported [40,44,53–57]. However, information about the prediction and selection of exclusive epitopes in ZIKV structural and nonstructural proteins has not yet been published. In this study, we used several bioinformatic tools to identify immunoreactive epitopes within ZIKV structural and nonstructural proteins and to see that they have no similarity with epitopes from other important flaviviruses. Therefore, these epitopes could be used to develop new vaccines and diagnostic tests for ZIKV.

The cross-reactivity of some antibodies is related to the high similarity between the amino acids and the tridimensional structure between epitopes [36]. The identification of exclusive epitopes for the development of ZIKV vaccines may contribute to avoid the ADE observed in some flaviviruses infections. The ADE predicts that previous infection with a flaviviruses could contribute to virus propagation in a subsequent infection by another related flaviviruses [58–60]. Although there are no definitive data that indicate that the ADE in humans, the overlap of *Flavivirus*-endemic areas [61] makes the ADE an important challenge in the elaboration of a vaccine candidate against ZIKV [62]. In this sense, it is reasonable to propose a vaccine platform based on epitopes to ZIKV that could induce an effective immunologic response without any cross-reacting activity against other flaviviruses.

Using *in silico* analysis, several groups have predicted ZIKV-specific mouse or human T-cell epitopes within E [40,44,54,63], NS2a [63], NS5 proteins [56] or the entire ZIKV polyprotein [45,55,64–67]. In this study, we found 91 epitopes with immunogenic properties on structural and nonstructural ZIKV proteins. Twenty-four epitopes had previously been identified (Supplementary Table S7). These studies showed that some epitopes (pr<sup>47–55</sup>, M<sup>56–70</sup>, E<sup>46–54</sup>, NS2a<sup>43–51</sup> and NS5<sup>19–27</sup>) were able to induce T-helper I type cytokines, including interferon gamma and/or tumor necrosis factor alpha secretion by CD4<sup>+</sup> and CD8<sup>+</sup> T cells in mice [63–65,67].

The data we present here show that the predicted epitopes for HLA-II binding are likely to bind to more alleles than the predicted epitopes for HLA-I binding. This could be related to the structure of the HLA-I and HLA-II molecules. The open structure of the binding groove and the longer length of HLA-II-bound peptides allow greater flexibility in binding and association with more HLA-II alleles [68]. Therefore, this could be responsible for the greater population coverage showed by predicted binding HLA-II epitopes (Fig. 5). It also identified a high number of predicted epitopes on the NS5 protein (384/26.36%). This result is probably the result of the length and immunogenicity of this protein [69–71]. The NS5 protein is the most conserved nonstructural protein of flaviviruses [72,73]. Therefore, because of the high conservation rate between the NS5 sequences of flaviviruses, only ten unique epitopes were identified on ZIKV NS5 protein (Supplementary Table S6).

Determination of the SAS is important to identify candidate epitopes exposed in the structural proteins and the NSI nonstructural protein. Exposed epitopes are more readily recognized by antibodies, and they are probably potential candidates for vaccine and diagnostic test development [74,75]. Previous studies showed that the interaction area of broadly neutralizing antibodies to DENV epitopes on protein E ranges from 1050 to 1400 Å<sup>2</sup> [76]. Most of the epitopes identified in

**TABLE 4.** List of candidate T-cell epitopes predicted from ZIKV structural and nonstructural proteins

Protein	Position	Epitope	HLA class	Antigenic score	Conservation among ZIKV strains (%)	Population coverage (%)	SAS (Å <sup>2</sup> )	Reference
E	374	MMLELDPPF	I	1.1639	100	59.52	1.217.23	[40,44]
	477	LNTKNGSISLMCLAL	II	1.7899	99.46	92.7	1.252.47	This study
C	61	PSLGLINRWGVSQK	II	1.4993	99.98	74.37	—	This study
	103	RRGADTSVIVGLLL	II	1.1073	98.47	93.45	—	[45]
pr	36	IQIMDLGHMCDATMS	II	1.0307	99.59	79.11	—	This study
NS2a	90	LVSFIFRANWTPRES	II	1.6900	99.88	99.5	—	[45]
	174	LATCGGFMLLSLKGK	II	1.2330	99.80	99.3	—	This study
	179	GFMLLSLKGKGSVKK	II	1.0070	99.78	62.28	—	This study
	190	SVKKNLFPVMALGLT	II	1.3171	99.91	99.78	—	This study
	195	LPFVMALGLTAVRLV	II	1.3379	99.50	99.69	—	[45]
	200	ALGLTAVRLVDPIV	II	1.6769	99.55	97.2	—	[45]
NS3	175	PVECFEFSMLKKQL	II	1.1434	99.82	98.09	—	This study
NS4a	82	GFGMVTLGASAWLMW	II	1.0313	99.97	99.6	—	This study
	99	EIEPARIACVLIVVF	II	1.1164	100	94.92	—	[45]

HLA, human leukocyte antigen; SAS, solvent-accessible surface; ZIKV, Zika virus.

M, E and NS1 proteins have a larger exposed surface area (Table 3). Therefore, it is possible that antibodies produced after ZIKV infection recognize these epitopes.

An ideal candidate ZIKV vaccine will be able to induce virus-specific immune response to all ZIKV strains [77]. Our analysis showed 91 potentially immunogenic epitopes present in the structural and nonstructural ZIKV proteins. Thus, considering these peptides are highly conserved among the 684 strains analysed in this study (90.89–100%), they are likely to be used in a vaccine formulation to induce protective immunity against any ZIKV strain. In order to identify the most promising candidates, we listed the epitopes that have a greater number of desired characteristics (conservation rate ≥95% among the ZIKV strains; population coverage >50%; antigenic score ≥1.0 and SAS ≥1050 Å<sup>2</sup>). The threshold for this classification was defined on the basis of the highest scores obtained in this study and prioritizing epitopes with ≥95% conservation because strains with a similarity of >95% were previously able to produce neutralizing antibodies for infections by different strains [78]. Among the epitopes that are able to bind HLA-I, E<sup>374–382</sup> is the only one that meets all the predetermined requirements, confirming the immunogenic potential of this epitope previously reported [40,44]. Thirteen HLA-II epitopes met all the requirements, including four epitopes present in structural proteins (pr<sup>36–50</sup>, C<sup>61–75</sup>, C<sup>103–117</sup>, E<sup>477–491</sup>) and 9 in nonstructural proteins (NS2a<sup>90–104</sup>, NS2a<sup>174–188</sup>, NS2a<sup>179–193</sup>, NS2a<sup>190–204</sup>, NS2a<sup>195–209</sup>, NS2a<sup>200–214</sup>, NS3<sup>175–189</sup>, NS4a<sup>82–96</sup>, NS4a<sup>99–113</sup>). Among the epitopes present in structural proteins, E<sup>477–491</sup> has the highest antigenic score (1.7899). This epitope is located in domain III of the E protein, which is the main domain associated with the induction of neutralizing antibodies [79,80]. Of the nine HLA-II epitopes present in the nonstructural proteins, six (66.66%) were identified on NS2a protein (Table 4). The NS2a protein is involved in viral RNA replication, virus assembly and secretion in flaviviruses [81]. It

was also demonstrated that ZIKV NS2a protein disrupts mammalian cortical neurogenesis by degrading adherens junction protein [82]. Therefore, epitopes predicted in this protein appear as potential candidates for the development of new vaccine formulations. Epitopes in NS2a have already been identified [45,63]. However, here we identify new and exclusive epitopes on the ZIKV NS2a protein (Table 4).

The 14 predicted T-cell epitopes identified in this study, taken together, are of special interest as potential candidate regions for inclusion in developing epitope-driven vaccines against ZIKV or diagnostic tests for the disease. Therefore, these epitopes have a biotechnologic relevance to the rational design of vaccine and diagnostic tests for ZIKV. However, in the next years, future *in vitro* and *in vivo* studies should be carried out to best characterize the immunogenicity of these epitopes and to establish the role of these epitopes as new vaccines or diagnostic tests.

### Acknowledgements

We thank S. Weaver (University of Texas Medical Branch) for his careful review. Supported in part by the FAPEMIG (grants APQ-01165–16 and PPM-00399–18). EAS received fellowships from FAPEMIG (PIBICT) and CNPq (PIBIC). GAPS received fellowships from FAPEMIG. LFLC received PQ fellowship from CNPq. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brasil (CAPES), Finance Code 001.

### Conflict of interest

None declared.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.nmni.2019.01.002>.

## References

- [1] Göertz GP, Abbo SR, Fros JJ, Pijlman GP. Functional RNA during *Zika virus* infection. *Virus Res* 2018;254:41–53.
- [2] Rückert C, Weger-Lucarelli J, Garcia-Luna SM, Young MC, Byas AD, Murrieta RA, et al. Impact of simultaneous exposure to arboviruses on infection and transmission by *Aedes aegypti* mosquitoes. *Nat Commun* 2017;8:15412.
- [3] Simmonds P, Becher P, Bukh J, Gould EA, Meyers G, Monath T, et al. ICTV virus taxonomy profile: *Flaviviridae*. *J Gen Virol* 2017;98:2–3.
- [4] Hou W, Cruz-cosme R, Armstrong N, Obwolo LA, Wen F, Hu W, et al. Molecular cloning and characterization of the genes encoding the proteins of *Zika virus*. *Gene* 2017;628:117–28.
- [5] Dick GW, Kitchen SF, Haddow AJ. *Zika virus*. I. Isolations and serological specificity. *Trans R Soc Trop Med Hyg* 1952;46:509–20.
- [6] Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. *Zika virus* outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med* 2009;360:2536–43.
- [7] Zinszer K, Morrison K, Brownstein JS, Marinho F, Santos AF, Nsoesie EO. Reconstruction of *Zika virus* introduction in Brazil. *Emerg Infect Dis* 2017;23:91–4.
- [8] Passos SRL, Borges dos Santos MA, Cerbino-Neto J, Buonora SN, Souza TML, de Oliveira RVC, et al. Detection of *Zika virus* in April 2013 patient samples, Rio de Janeiro, Brazil. *Emerg Infect Dis* 2017;23:2120–1.
- [9] Musso D. *Zika virus* transmission from French Polynesia to Brazil. *Emerg Infect Dis* 2015;21:1887–9.
- [10] Oehler E, Watrin L, Larre P, Leparc-Goffart I, Lastere S, Valour F, et al. *Zika virus* infection complicated by Guillain-Barre syndrome—case report, French Polynesia, December 2013. *Euro Surveill* 2014;19:20720.
- [11] Edge L. *Zika virus* on the move. *Cell* 2016;164:585–7.
- [12] Malkki H. *Zika virus* infection could trigger Guillain-Barré syndrome. *Nat Rev Neurol* 2016;12:187.
- [13] Musso D, Roche C, Robin E, Nhan T, Teissier A, Cao-Lormeau VM. Potential sexual transmission of *Zika virus*. *Emerg Infect Dis* 2015;21:359–61.
- [14] Zamani M, Zamani V. Sexual transmission of *Zika virus*: an assessment of the evidence. *Iran J Public Health* 2017;46:1305–6.
- [15] Tebas P, Roberts CC, Muthumani K, Reuschel EL, Kudchodkar SB, Zaidi FI, et al. Safety and immunogenicity of an anti-Zika virus DNA vaccine – preliminary report. *N Engl J Med* 2017. <https://doi.org/10.1056/NEJMoa1708120>.
- [16] Gaudinski MR, Houser KV, Morabito KM, Hu Z, Yamshchikov G, Rothwell RS, et al. Safety, tolerability, and immunogenicity of two *Zika virus* DNA vaccine candidates in healthy adults: randomised, open-label, phase I clinical trials. *Lancet* 2018;391(10120):552–62.
- [17] Richner JM, Himansu S, Dowd KA, Pierson TC, Ciaramella G, Diamond MS, et al. Modified mRNA vaccines protect against *Zika virus* infection. *Cell* 2017;168. 1114–25.e10.
- [18] Barouch DH, Thomas SJ, Michael NL. Perspective prospects for a *Zika virus* vaccine. *Immunity* 2017;46:176–82.
- [19] Barrett ADT. Current status of *Zika* vaccine development: *Zika* vaccines advance into clinical evaluation. *NPJ Vaccines* 2018;3:24.
- [20] Culshaw A, Mongkolsapaya J, Srean G. The immunology of *Zika virus*. *Fl000Res* 2018;7:203.
- [21] Kawiecki AB, Christofferson RC. *Zika virus*–induced antibody response enhances dengue virus serotype 2 replication *in vitro*. *J Infect Dis* 2016;214:1357–60.
- [22] Priyamvada L, Quicke KM, Hudson WH, Onlamoon N, Sewatanon J, Edupuganti S, et al. Human antibody responses after dengue virus infection are highly cross-reactive to *Zika virus*. *Proc Natl Acad Sci U S A* 2016;113:7852–7.
- [23] Rey FA, Stiasny K, Vaney M, Dellarole M, Heinz FX. The bright and the dark side of human antibody responses to flaviviruses: lessons for vaccine design. *EMBO Rep* 2018;19:206–24.
- [24] Chiang MH, Sung WC, Lien SP, Chen YZ, Lo AF, Huang JH, et al. Identification of novel vaccine candidates against *Acinetobacter baumannii* using reverse vaccinology. *Hum Vaccin Immunother* 2015;11:1065–73.
- [25] Contreras M, Villar M, Artigas-Jerónimo S, Kornieieva L, Mytrofanov S, de la Fuente J. A reverse vaccinology approach to the identification and characterization of *Ctenocephalides felis* candidate protective antigens for the control of cat flea infestations. *Parasit Vectors* 2018;11:43.
- [26] Backert L, Kohlbacher O. Immunoinformatics and epitope prediction in the age of genomic medicine. *Genome Med* 2015;7:119.
- [27] Bruno L, Cortese M, Rappuoli R, Merola M. Lessons from reverse vaccinology for viral vaccine design. *Curr Opin Virol* 2015;11:89–97.
- [28] Heinson AI, Woelk CH, Newell ML. The promise of reverse vaccinology. *Int Health* 2015;7:85–9.
- [29] Liljeroos L, Malito E, Ferlenghi I, Bottomley MJ. Structural and computational biology in the design of immunogenic vaccine antigens. *J Immunol Res* 2015;2015:1–17.
- [30] Schussek S, Trieu A, Doolan DL. Genome- and proteome-wide screening strategies for antigen discovery and immunogen design. *Biotechnol Adv* 2014;32:403–14.
- [31] Huber SR, van Beek J, de Jonge J, Luytjes W, van Baarle D. T cell responses to viral infections—opportunities for peptide vaccination. *Front Immunol* 2014;5(APR):1–12.
- [32] Nandy A, Basak SC. A brief review of computer-assisted approaches to rational design of peptide vaccines. *Int J Mol Sci* 2016;17:666.
- [33] Patronov A, Doytchinova I. T-cell epitope vaccine design by immunoinformatics. *Open Biol* 2013;3:120139.
- [34] Paul S, Sidney J, Sette A, Peters B. TepiTool: a pipeline for computational prediction of T cell epitope candidates. *Curr Protoc Immunol* 2016;18.19.1–18.19.24.
- [35] Bui HH, Sidney J, Li W, Füsseder N, Sette A. Development of an epitope conservancy analysis tool to facilitate the design of epitope-based diagnostics and vaccines. *BMC Bioinform* 2007;8:361.
- [36] Breiteneder H, Mills C. Structural bioinformatic approaches to understand cross-reactivity. *Mol Nutr Food Res* 2006;50:628–32.
- [37] Negi SS, Braun W. Cross-React: a new structural bioinformatics method for predicting allergen cross-reactivity. *Bioinformatics* 2017;33:1014–20.
- [38] Doytchinova IA, Flower DR. Vaxijen: a server for prediction of protective antigens, tumour antigens and subunit vaccines. *BMC Bioinform* 2007;8:4.
- [39] Gupta S, Kapoor P, Chaudhary K, Gautam A, Kumar R. Open Source Drug Discovery Consortium, et al. *In silico* approach for predicting toxicity of peptides and proteins. *PLoS One* 2013;8:e73957.
- [40] Janahi EM, Dhasmana A, Srivastava V, Sarangi AN, Raza S, Arif JM, et al. *In silico* CD4<sup>+</sup>, CD8<sup>+</sup> T-cell and B-cell immunity associated immunogenic epitope prediction and HLA distribution analysis of *Zika virus*. *EXCLI J* 2017;16:63–72.
- [41] Solanki V, Tiwari V. Subtractive proteomics to identify novel drug targets and reverse vaccinology for the development of chimeric vaccine against *Acinetobacter baumannii*. *Sci Rep* 2018;8:9044.
- [42] Zheng J, Lin X, Wang X, Zheng L, Lan S, Jin S, et al. *In silico* analysis of epitope-based vaccine candidates against hepatitis B virus polymerase protein. *Viruses* 2017;9:112.

- [43] Bui HH, Sidney J, Dinh K, Southwood S, Newman MJ, Sette A. Predicting population coverage of T-cell epitope-based diagnostics and vaccines. *BMC Bioinform* 2006;7:153.
- [44] Alam A, Ali S, Ahmad S, Malik MZ, Ishrat R. From ZikV genome to vaccine: *in silico* approach for the epitope-based peptide vaccine against Zika virus envelope glycoprotein. *Immunology* 2016;149:386–99.
- [45] Ramaiah A, Dai L, Contreras D, Sinha S, Sun R, Arumugaswami V. Comparative analysis of protein evolution in the genome of pre-epidemic and epidemic Zika virus. *Infect Genet Evol* 2017;51:74–85.
- [46] Marano G, Pupella S, Vaglio S, Liunbruno GM, Grazzini G. Zika virus and the never-ending story of emerging pathogens and transfusion medicine. *Blood Transfus* 2016;14:95–100.
- [47] Beckham JD, Pastula DM, Massey A, Tyler KL. Zika virus as an emerging global pathogen: neurological complications of Zika virus. *JAMA Neurol* 2016;73:875–9.
- [48] Basarab M, Bowman C, Aarons EJ, Cropley I. Zika virus. *BMJ* 2016;352:1–7.
- [49] Makhluf H, Shrestha S. Development of Zika virus vaccines. *Vaccines* 2018;6:7.
- [50] Du L, Zhou Y, Jiang S. The latest advancements in Zika virus vaccine development. *Expert Rev Vaccines* 2017;16:951–4.
- [51] Usman Mirza M, Rafique S, Ali A, Munir M, Ikram N, Manan A, et al. Towards peptide vaccines against Zika virus: immunoinformatics combined with molecular dynamics simulations to predict antigenic epitopes of Zika viral proteins. *Sci Rep* 2016;6:37313.
- [52] Noor R, Ahmed T. Zika virus: epidemiological study and its association with public health risk. *J Infect Public Health* 2018;11:611–6.
- [53] Yasmin T, Akter S, Debnath M, Ebihara A, Nakagawa T, Nabi AHMN. *In silico* proposition to predict cluster of B- and T-cell epitopes for the usefulness of vaccine design from invasive, virulent and membrane associated proteins of *C. jejuni*. *Silico Pharmacol* 2016;4:5.
- [54] Ashfaq UA, Ahmed B. De novo structural modeling and conserved epitopes prediction of Zika virus envelop protein for vaccine development. *Viral Immunol* 2016;29:436–43.
- [55] Dar H, Zaheer T, Rehman MT, Ali A, Javed A, Khan GA, et al. Prediction of promiscuous T-cell epitopes in the Zika virus polyprotein: an *in silico* approach. *Asian Pac J Trop Med* 2016;9:844–50.
- [56] dos Santos Franco L, Oliveira Vidal P, Amorim JH. *In silico* design of a Zika virus non-structural protein 5 aiming vaccine protection against Zika and dengue in different human populations. *J Biomed Sci* 2017;24:88.
- [57] Ebrahimi S, Mohabatkari H. Prediction of T-cell epitopes for designing a reverse vaccine against streptococcal bacteria. *Mol Biol Res Commun* 2018;7:35–41.
- [58] Dejnirattisai W, Supasa P, Wongwiwat W, Rouvinski A, Barba-Spaeth G, Duangchinda T, et al. Dengue virus sero-cross-reactivity drives antibody-dependent enhancement of infection with Zika virus. *Nat Immunol* 2016;17:1102–8.
- [59] Kam YW, Lee CYP, Teo TH, Howland SW, Amrun SN, Lum FM, et al. Cross-reactive dengue human monoclonal antibody prevents severe pathologies and death from Zika virus infections. *JCI Insight* 2017;2:e92428.
- [60] Paul LM, Carlin ER, Jenkins MM, Tan AL, Barcellona CM, Nicholson CO, et al. Dengue virus antibodies enhance Zika virus infection. *Clin Transl Immunol* 2016;5:e117.
- [61] Azeredo EL, Dos Santos FB, Barbosa LS, Souza TMA, Badolatto-Correa J, Sanchez-Arcila JC, et al. Clinical and laboratory profile of Zika and dengue infected patients: lessons learned from the co-circulation of dengue, Zika and chikungunya in Brazil. *PLoS Curr* 2018;10.
- [62] Fernandez E, Diamond MS. Vaccination strategies against Zika virus. *Curr Opin Virol* 2017;23:59–67.
- [63] Huang H, Li S, Zhang Y, Han X, Jia B, Liu H, et al. CD8(+) T cell immune response in immunocompetent mice during Zika virus infection. *J Virol* 2017;91:e00900–17.
- [64] Hassert M, Wolf KJ, Schwetye KE, DiPaolo RJ, Brien JD, Pinto AK. CD4<sup>+</sup> T cells mediate protection against Zika associated severe disease in a mouse model of infection. *PLoS Pathog* 2018;14:e1007237.
- [65] Ngono AE, Vizcarra EA, Tang WW, Sheets N, Joo Y, Kim K, et al. Mapping and role of the CD8(+) T cell response during primary Zika virus infection in mice. *Cell Host Microbe* 2017;21:35–46.
- [66] Dikhit MR, Ansari Y, Mansuri R, Sahoo BR, Dehury B, Amit A, et al. Infection, genetics and evolution computational prediction and analysis of potential antigenic CTL epitopes in Zika virus: a first step towards vaccine development. *MEEGID* 2016;45:187–97.
- [67] Wen J, Tang WW, Sheets N, Ellison J, Sette A, Kim K, et al. Identification of Zika virus epitopes reveals immunodominant and protective roles for dengue virus cross-reactive CD8<sup>+</sup> T cells. *Nat Microbiol* 2017;2:17036.
- [68] Wiczorek M, Abualrous ET, Sticht J, Álvaro-Benito M, Stolzenberg S, Noé F, et al. Major histocompatibility complex (MHC) class I and MHC class II proteins: conformational plasticity in antigen presentation. *Front Immunol* 2017;8:292.
- [69] Laurent-Rolle M, Morrison J, Rajsbaum R, Macleod JML, Pisanelli G, Pham A, et al. The interferon signaling antagonist function of yellow fever virus NS5 protein is activated by type I interferon. *Cell Host Microbe* 2014;16:314–27.
- [70] Morrison J, Laurent-Rolle M, Maestre AM, Rajsbaum R, Pisanelli G, Simon V, et al. Dengue virus co-opts UBR4 to degrade STAT2 and antagonize type I interferon signaling. *PLoS Pathog* 2013;9:e1003265.
- [71] Lubick KJ, Robertson SJ, McNally KL, Boehm M, Yoshii K, Best SM, et al. Flavivirus antagonism of type I interferon signaling reveals prolidase as a regulator of IFNAR1 surface expression. *Cell Host Microbe* 2015;18:61–74.
- [72] Grant A, Ponia SS, Tripathi S, Balasubramanian V, Miorin L, Sourisseau M, et al. Zika virus targets human STAT2 to inhibit type I interferon signaling. *Cell Host Microbe* 2016;19:882–90.
- [73] Iacono-Connors LC, Schmaljohn CS. Cloning and sequence analysis of the genes encoding the nonstructural proteins of langat virus and comparative analysis with other flaviviruses. *Virology* 1992;188:875–80.
- [74] Dowd KA, Mukherjee S, Kuhn RJ, Pierson TC. Combined effects of the structural heterogeneity and dynamics of flaviviruses on antibody recognition. *J Virol* 2014;88:11726–37.
- [75] Heinz FX, Stiasny K. Flaviviruses and their antigenic structure. *J Clin Virol* 2012;55:289–95.
- [76] Rouvinski A, Guardado-Calvo P, Barba-Spaeth G, Duquerroy S, Vanev MC, Kikuti CM, et al. Recognition determinants of broadly neutralizing human antibodies against dengue viruses. *Nature* 2015;520(7545):109–13.
- [77] Beaver JT, Lelutiu N, Habib R, Skountzou I. Evolution of two major Zika virus lineages: implications for pathology, immune response, and vaccine development. *Front Immunol* 2018;9:1640.
- [78] Dowd KA, DeMaso CR, Pelc RS, Speer SD, Smith ARY, Goo L, et al. Broadly neutralizing activity of Zika virus-immune sera identifies a single viral serotype. *Cell Rep* 2016;16:1485–91.
- [79] Deng YQ, Dai JX, Ji GH, Jiang T, Wang HJ, Yang H, et al. A broadly Flavivirus cross-neutralizing monoclonal antibody that recognizes a novel epitope within the fusion loop of E protein. *PLoS One* 2011;6:e16059.
- [80] Chávez JH, Silva JR, Amarilla AA, Moraes Figueiredo LT. Domain III peptides from Flavivirus envelope protein are useful antigens for serologic diagnosis and targets for immunization. *Biologicals* 2010;38:613–8.
- [81] Leung JY, Pijlman GP, Kondratieva N, Hyde J, Mackenzie JM, Khromykh AA. Role of nonstructural protein NS2A in Flavivirus assembly. *J Virol* 2008;15(82):4731–41.
- [82] Yoon KJ, Song G, Qian X, Pan J, Xu D, Rho HS, et al. Zika virus-encoded NS2A disrupts mammalian cortical neurogenesis by degrading adherens junction proteins. *Cell Stem Cell* 2017;21:349–358.e6.